

Exponential cosmological redshift in a linearly expanding universe

Neil V. Budko

*Laboratory of Electromagnetic Research, Faculty of Electrical Engineering, Mathematics and Computer Science,
Delft University of Technology, Mekelweg 4, 2628 CD Delft, The Netherlands**

(Dated: July 3, 2009)

The first principles analysis of the radiation by an arbitrary source in a flat Friedmann-Robertson-Walker space-time is presented. The obtained analytical solution explicitly shows that the cosmological redshift is not of kinematic origin and that the source and the observer may be regarded as being at rest with respect to each other at all times. At the same time the effect of the time-variation of the metric on the propagation of light appears to be underestimated in the standard cosmology. The cosmological redshift caused by the linear time-variation of the metric turns out to be an exponential rather than linear function of the well-defined spatial distance and the apparent brightness of the source contains an even stronger exponential decay factor.

PACS numbers: 03.50.De, 03.65.Pm, 04.20.Jb

The distance-dependent redshift of the electromagnetic radiation from stellar sources is the key observational parameter in astronomy and a reality-check for any viable cosmological theory. The expanding universe model described in general relativity by the Friedmann-Robertson-Walker (FRW) metric [1] is accepted as the most natural explanation of the observed data, i.e., the Hubble's approximately linear law. This law has been deduced experimentally by estimating the distances to particular types of redshifted stellar sources – standard astronomical candles – from the measurement of their apparent brightness [2]. The preferred theoretical explanation goes back to Lemaitre [3] and is based on the analysis of the light-cone condition for the space-time interval in the FRW metric. However, the original *linear* Hubble's redshift law is recovered only in the “small” distance (small redshift) approximation [4]. An exact law universally valid for large distances and large redshifts has never been derived. Recently we have seen a surge in the astronomical data containing extremely high redshifts [5]. There is also a growing interest in various new cosmological scenarios, e.g. inflation, accelerated expansion, varying speed of light theories, and cyclic universes [6]. Since all of these theories and data need to be tested against the simplest flat-space linearly expanding universe model, the exact redshift and apparent brightness (or luminosity) formulas resulting from this basic cosmological model and valid for all distances have become especially important.

On the other hand, the physical meaning of the expanding universe model is also still open to debate [7]. What exactly is expanding? Does the spatial separation between galaxies actually grow in time? How can this mutual recession be superluminal (for higher redshifts)? These elementary questions seem to have as many answers as there are cosmologists, with the majority favouring a more or less literal picture of galaxies flying apart. This is often appended by a vague statement that they are not flying apart *in* space, but rather

because the space itself is expanding. This interpretation is an important part of the *standard model of cosmology*. In particular, the literal expansion interpretation means that in the past all objects were physically closer to each other, and that there was a very densely packed state in the beginning – the Big Bang.

Being a very simple algebraic expression, the space-time interval, which forms the backbone of the modern cosmological analysis, unfortunately, allows conflicting interpretations still causing many confusions and heated debates [7]. Even the most basic conclusions are not as clearcut as it seems. For example, being shown the interval in the flat expanding FRW metric

$$ds^2 = c^2 dt^2 - a^2(t) (dx_1^2 + dx_2^2 + dx_3^2), \quad (1)$$

we usually hear something like this: The change of the scale factor $a(t)$ with time, e.g. linear growth, means that all spatial distances are increasing with time. Hence, all objects will be receding from each other. This seemingly obvious statement, in fact, does not follow from (1) at all. What we see in (1) is that the spatial interval is changing with respect to the *time-interval*. Whether this means an increase of the spatial distance with *time* as well remains to be seen. In particular, does the *co-moving* distance $a(t)R$ have any physical meaning?

Here and in the companion paper [8] devoted to the wavepropagation in non-autonomous time-varying media I describe a general first-principles approach which may provide a definitive resolution of this and other interpretational problems. I obtain the exact redshift-distance relation valid for all distances and show the way to derive the exact brightness (luminosity) formula. The basic idea goes back to Schrödinger and amounts to first posing and solving a general-relativistic problem for an arbitrary electromagnetic source radiating into a homogeneous infinite background with an FRW metric, and only then trying to find the physical interpretation. Fortunately, the solution is analytical and its complete derivation can be found in [8]. However, it does have several surpris-

ing features, which in my view call for a second look at our redshift data and may even require a revision of the standard interpretation mentioned above. First of all, the solution explicitly shows that the cosmological redshift is not of kinematic origin. Secondly, the influence of a time-varying metric on the electromagnetic field seems to be severely underestimated in the standard cosmology. Not only the exact Hubble law turns out to be exponential rather than linear, but there is also a very strong exponential decay in the apparent brightness of the source. The latter is heavily relied upon in the verification of the Hubble law and to find possible deviations from its linear form for higher redshifts.

The earliest and, probably, the most complete attempt along the present lines is the Schrödinger's 1939 paper [9]. Schrödinger, however, considered the covariant *scalar* wave equation without the source term, which, strictly speaking, is not the equation governing the electromagnetic radiation by a causal source. The first-principles analysis should start with the covariant first-order Maxwell's system, and proceed by *deriving* the corresponding wave equation, including the source-term, and solving it. Although, the first-order formulation of the problem has been considered many times since [10, 11, 12], no deviations from the original linear Hubble's law were reported.

In Cartesian coordinates the FRW metric of a spatially flat expanding universe is

$$[g_{\alpha\beta}] = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & -a^2(t) & 0 & 0 \\ 0 & 0 & -a^2(t) & 0 \\ 0 & 0 & 0 & -a^2(t) \end{bmatrix}, \quad (2)$$

with determinant $g = -a^6(t)$, where $a(t)$ is the curvature radius determining the expansion rate. As was demonstrated by Tamm [10, 13, 14], the electromagnetic field in an empty, but curved space-time, at each location $\mathbf{x}$ and with respect to the local time t satisfies the usual Maxwell's equations in the $(3+1)$ -form

$$\begin{aligned} -\nabla \times \mathbf{H}(\mathbf{x}, t) + \partial_t \mathbf{D}(\mathbf{x}, t) &= -\sqrt{-g} \mathbf{J}(\mathbf{x}, t), \\ \nabla \times \mathbf{E}(\mathbf{x}, t) + \partial_t \mathbf{B}(\mathbf{x}, t) &= 0, \end{aligned} \quad (3)$$

with the following metric-induced local constitutive relations:

$$\begin{aligned} \mathbf{D}(\mathbf{x}, t) &= \varepsilon_0 \varepsilon \mathbf{E}(\mathbf{x}, t) + \frac{1}{c} \mathbf{w} \times \mathbf{H}(\mathbf{x}, t), \\ \mathbf{B}(\mathbf{x}, t) &= \mu_0 \mu \mathbf{H}(\mathbf{x}, t) - \frac{1}{c} \mathbf{w} \times \mathbf{E}(\mathbf{x}, t), \end{aligned} \quad (4)$$

where

$$\varepsilon = \mu = -\frac{\sqrt{-g}}{g_{00}} g^{ij}, \quad w_i = \frac{g_{0i}}{g_{00}}, \quad (5)$$

with $i, j = 1, 2, 3$, and $[g^{\alpha\mu}] = [g_{\alpha\mu}]^{-1}$. For the FRW metric (2) these relations reduce to

$$\begin{aligned} \mathbf{D}(\mathbf{x}, t) &= \varepsilon_0 a(t) \mathbf{E}(\mathbf{x}, t), \\ \mathbf{B}(\mathbf{x}, t) &= \mu_0 a(t) \mathbf{H}(\mathbf{x}, t), \end{aligned} \quad (6)$$

Substituting these into (3) we arrive at the following Maxwell's equations:

$$\begin{aligned} -\nabla \times \mathbf{B}(\mathbf{x}, t) + \mu_0 a(t) \partial_t \mathbf{D}(\mathbf{x}, t) &= -\mu_0 a^4(t) \mathbf{J}(\mathbf{x}, t), \\ \nabla \times \mathbf{D}(\mathbf{x}, t) + \varepsilon_0 a(t) \partial_t \mathbf{B}(\mathbf{x}, t) &= 0. \end{aligned} \quad (7)$$

Let us clarify the physical meaning of our coordinates. The Maxwell equations (7) describe the *local* behaviour of the electromagnetic field. The time-variable involved is called the *synchronous time* [4], meaning that any physical process, say, emission of N photons, which takes T -seconds of local t -time at some spatial location will take exactly T -seconds of local t -time, if it would happen at another location. The fact that we can write the covariant Maxwell's equations in the simple form (7) and not worry about the choice of the spatial coordinate system is due to the spatial flatness of the FRW metric, which we assumed at the beginning. Any (flat, stationary, no stretching, etc.) rectangular Cartesian system would do [4].

Now, we introduce a variable change

$$\tau(t) = \int_{t_0}^t \frac{dt'}{a(t')}, \quad (8)$$

where t_0 is the “switch-on” moment of the causal source $\mathbf{J}(\mathbf{x}, t)$. The new time-like variable is known as the *cosmological time*, and the corresponding variable change procedure is sometimes called a *conformal transformation* [7], since it transforms the general-relativistic FRW metric into the Minkowski metric of special relativity. Indeed, in terms of this new variable the Maxwell equations become

$$\begin{aligned} -\nabla \times \mathbf{B}(\mathbf{x}, \tau) + \mu_0 \partial_\tau \mathbf{D}(\mathbf{x}, \tau) &= -\mu_0 a^4(\tau) \mathbf{J}(\mathbf{x}, \tau), \\ \nabla \times \mathbf{D}(\mathbf{x}, \tau) + \varepsilon_0 \partial_\tau \mathbf{B}(\mathbf{x}, \tau) &= 0, \end{aligned} \quad (9)$$

and are the same as Maxwell's equations in Minkowski vacuum up to the right-hand side and the above variable change. At this moment we still cannot say whether the source is receding with respect to the observation point or not, as the Maxwell equations represent a *local* relation between the current density and the fields at the same point in space. To make any conclusions about the possible physical recession we need an explicit solution of these equations relating the current density at one location to the fields at some other location.

From the mathematical point of view, the right-hand side of (7) contains functions of τ which must be interpreted as $a(t(\tau))$ and $\mathbf{J}(\mathbf{x}, t(\tau))$, where $t(\tau)$ is a function inverse with respect to $\tau(t)$. Obviously, this inverse function is well-defined whenever $a(t) > 0$ for all

$t \geq t_0$. Equations (9) are identical in form to the vacuum Maxwell's equations and therefore have a well-known analytical solution. In particular, they can be reduced [8] to the following scalar wave equation:

$$\Delta \mathbf{D}(\mathbf{x}, \tau) - \frac{1}{c_0^2} \partial_\tau^2 \mathbf{D}(\mathbf{x}, \tau) = \frac{1}{c_0^2} \partial_\tau [a^4(\tau) \mathbf{J}(\mathbf{x}, \tau)] - \nabla(\nabla \cdot) \int_0^\tau a^4(\tau') \mathbf{J}(\mathbf{x}, \tau') d\tau'. \quad (10)$$

Now we can see the crucial difference between the present approach and that of Schrödinger [9]. Although the left-hand side of (10) is formally the same as the one obtained in [9], Schrödinger had to use a different variable change to arrive at it. Indeed, the covariant approach to a scalar wave equation would have resulted in $a^3(t)$ in the denominator of (8), whereas what we have used here is in perfect agreement with the usual definition of the cosmological time, represents a conformal transformation, and appears in other first-order formulations of the problem [10, 11, 12]. The explicit solution of (10) in $(\mathbf{x}, \tau)$ -domain is:

$$\begin{aligned} \mathbf{E}(\mathbf{x}, \tau) = & \int_{\mathbf{x}' \in \mathbb{R}^3} \frac{[3\mathbb{Q} - \mathbb{I}]}{4\pi|\mathbf{x} - \mathbf{x}'|^3} \int_0^{\tau_r} \frac{a^4(\tau') \mathbf{J}(\mathbf{x}', \tau')}{\varepsilon_0 a(\tau)} d\tau' d\mathbf{x}' \\ & + \int_{\mathbf{x}' \in \mathbb{R}^3} \frac{[3\mathbb{Q} - \mathbb{I}]}{4\pi|\mathbf{x} - \mathbf{x}'|^2} \frac{a^4(\tau_r) \mathbf{J}(\mathbf{x}', \tau_r)}{c_0 \varepsilon_0 a(\tau)} d\mathbf{x}' \\ & + \int_{\mathbf{x}' \in \mathbb{R}^3} \frac{[\mathbb{Q} - \mathbb{I}]}{4\pi|\mathbf{x} - \mathbf{x}'|} \frac{\partial_\tau [a^4(\tau) \mathbf{J}(\mathbf{x}', \tau)]|_{\tau=\tau_r}}{c_0^2 \varepsilon_0 a(\tau)} d\mathbf{x}', \end{aligned} \quad (11)$$

In this formula

$$[\mathbb{Q}(\mathbf{x} - \mathbf{x}')]_{ij} = \frac{(x_i - x'_i)(x_j - x'_j)}{|\mathbf{x} - \mathbf{x}'|^2}, \quad (12)$$

and $\mathbb{I}$ denotes the Kronecker (3×3) -identity tensor. The retarded τ -time is

$$\tau_r = \tau - \frac{|\mathbf{x} - \mathbf{x}'|}{c_0}. \quad (13)$$

Apart from the source modulation (discussed below), the obtained $(\mathbf{x}, \tau)$ -domain solution is mathematically identical to the $(\mathbf{x}, t)$ -domain free-space radiation formula [8]. It does not provide any clear evidence of the mutual recession between the source at $\mathbf{x}'$ and the observer at $\mathbf{x}$. In fact, there are only two possibilities for the recession to enter our solution: either in the form of the co-moving distance $a(\tau)|\mathbf{x} - \mathbf{x}'|$ in terms of the cosmological time τ , or in terms of the synchronous time as $a(t)|\mathbf{x} - \mathbf{x}'|$. If the recession is really happening, then this co-moving distance should appear everywhere, where one would normally have a stationary distance $|\mathbf{x} - \mathbf{x}'|$, i.e. both in

the distance-dependent decay factors and in the retarded time expression. At the moment we can already dismiss the first of the above possibilities. Of course, we could artificially group the $a(\tau)$ terms in the denominator with the corresponding $|\mathbf{x} - \mathbf{x}'|$ terms. However, as shown below, this is mainly influencing the magnitude of the received field. For the recession to really *cause* the redshift the co-moving distance should be present in the temporal argument of the current density, i.e. in the retarded τ -time (13), which is obviously not the case.

If there is no mutual recession, where does the redshift come from? Mathematically, the reason for the cosmological redshift may be formulated as follows. The (electromagnetic) processes that are synchronous in the local synchronous time t are not synchronous in the cosmological time τ , and vice versa. To show this explicitly we need to re-write the $(\mathbf{x}, \tau)$ -domain radiation formula (11) as an $(\mathbf{x}, t)$ -domain formula, which requires an inverse transformation $t(\tau)$. For a general expansion rate $a(t) > 0$ this inverse transformation exists but is implicit. For the linear case, $a(t) = a_0 + bt$, however, the integral (8) can be evaluated and the resulting algebraic equation can be solved for t as:

$$t = \frac{a_0 + bt_0}{b} e^{b\tau} - \frac{a_0}{b}, \quad (14)$$

which also means

$$a(\tau) = (a_0 + bt_0) e^{b\tau}. \quad (15)$$

The most obvious demonstration of the redshift comes from considering a monochromatic source [8]. In that case the right-hand side of (11) contains trigonometric functions, say $\sin(\omega t)$. In view of (14) and (13) this functions will enter (11) as

$$\sin(\omega t(\tau_r)) = \sin \left(\omega \left(\frac{a_0 + bt_0}{b} e^{b(\tau - |\mathbf{x} - \mathbf{x}'|/c_0)} - \frac{a_0}{b} \right) \right), \quad (16)$$

or, recognizing here the t -domain result, as

$$\sin \left(\omega' t + \frac{a_0}{b} (\omega' - \omega) \right), \quad (17)$$

where the new frequency is

$$\omega' = \omega e^{-\frac{b}{c_0} |\mathbf{x} - \mathbf{x}'|}. \quad (18)$$

Hence, a monochromatic signal sent at frequency ω from $\mathbf{x}'$ will be observed at $\mathbf{x}$ at a lower frequency ω' . Moreover, this cosmological redshift will have an exact exponential dependence on the distance between the source and the observer. At this point we can dismiss the other possibility for the mutual recession between the source and the observer. We have just analyzed the retarded time expression in the $(\mathbf{x}, t)$ -domain and do not see any signs of the co-moving distance $a(t)|\mathbf{x} - \mathbf{x}'|$ in it. On the bright

side, the present interpretation is completely free from any conceptual problems as far as redshifts larger than one are concerned. Since their origin is clearly not kinematic, no superluminal velocities are required to explain them. It is also interesting to note that the redshift is completely independent of the initial moment of expansion, as long as this initial moment was before or exactly at t_0 – the moment at which the source starts radiating.

This redshift effect is a case of the general *time dilation*, which is confirmed by the observed apparent change in the duration of various well-understood astronomical processes. Let the two characteristic time instants in the evolution of the source current density be t_1^{src} and t_2^{src} . The local-time interval at the source location $\mathbf{x}'$ is then

$$\Delta t^{\text{src}} = t_1^{\text{src}} - t_2^{\text{src}} = \frac{a_0 + bt_0}{b} \left[e^{b\tau_1^{\text{src}}} - e^{b\tau_2^{\text{src}}} \right], \quad (19)$$

where τ_1^{src} and τ_2^{src} are the corresponding τ -times. To derive the corresponding local times at the observer location we apply (13) in (14) and obtain:

$$\Delta t^{\text{rec}} = t_1^{\text{rec}} - t_2^{\text{rec}} = \frac{a_0 + bt_0}{b} \left[e^{b\tau_1^{\text{src}}} - e^{b\tau_2^{\text{src}}} \right] e^{\frac{b}{c_0}|\mathbf{x}-\mathbf{x}'|}. \quad (20)$$

Hence, it follows that

$$\frac{\Delta t^{\text{rec}}}{\Delta t^{\text{src}}} = e^{\frac{b}{c_0}|\mathbf{x}-\mathbf{x}'|}. \quad (21)$$

Thus, the general time-dilation in a flat universe linearly expanding at a constant rate b is an exponential function of the distance between the source and the observer. In [8] the redshift formula for an exponentially expanding (de Sitter) universe is derived. Similarly to the linear case, it is a function of distance which mathematically does not even resemble the expansion rate function $a(t)$.

So far we have considered just one effect of the time-variation of the metric – the cosmological redshift. In addition to that the obtained analytical solution (11) shows the modulation of the source by the fourth power of the expansion rate. This demonstrates that there is more to the time-varying metric than a simple change of the rate of emission due to the time dilation. Substitution of (15) into (11) with the retarded time (13) will produce a multiplier of the form

$$a(\tau_r) = (a_0 + bt_0) e^{b\tau_r} = a(\tau) e^{-\frac{b}{c_0}|\mathbf{x}-\mathbf{x}'|}. \quad (22)$$

Since this term comes in the fourth power, the additional decay in the luminosity is by far stronger than would follow from the reduced emission rate. In fact, the sources are disappearing from our view significantly faster than they get redshifted. This means that the exact redshift-luminosity relation for a linearly expanding universe, which still needs to be worked out, is probably also different from the currently used one, especially at higher redshifts.

The modulation of the current produces another curious effect. We know that a DC-like current does not normally radiate. This follows from the fact that the far-field term for a vacuum background contains the time-derivative of the current. However, now the current is multiplied by the expansion rate. Thus, a DC-like current in a time-varying metric will radiate into the far-field zone. On the other hand, a time-varying current, which normally does radiate, will be effectively silenced, if its time variation happens to coincide with $a^{-4}(\tau)$.

In conclusion, the derived analytical radiation formula for an arbitrary source in an infinite, flat, linearly expanding universe provides the exact exponential redshift/time-dilation law valid for all distances and free from interpretational problems for higher redshifts. It appears that both the time-dilation and the decrease in the apparent brightness of the sources caused by the interaction of light with a time-varying FRW metric have so far been greatly underestimated. The obtained solution provides no convincing evidence for the change of the distances between the source and the observer in the FRW cosmology. Paradoxically, expanding universe does not follow from the expanding metric.

* Electronic address: n.v.budko@tudelft.nl

- [1] S. Weinberg, *Cosmology*, Oxford Univ. Press, USA, 2008.
- [2] E. Hubble, *Proc. NAS USA* **15**, 3, 168–173, 1929.
- [3] A. G. Lemaitre, *Mon. Not. R. Astron. Soc.* **91**, 483–490, 1927.
- [4] L. D. Landau and E. M. Lifshitz, *The Classical Theory of Fields*, Butterworth-Heinemann, 1980.
- [5] K. M. Lanzetta, A. Yahil, and A. Fernandez-Soto, *Nature* **381**, 759–763, 1996.
- [6] J. Khoury, P. J. Steinhardt, and N. Turok, *Phys. Rev. Lett.* **92**, 031302, 2004.
- [7] G. F. Ellis and W. Stoeger, *Class. Quant. Grav.* **5**, 207–220, 1988; L. B. Okun, K. G. Selivanov, and V. L. Telegdi, *Am. J. Phys.* **68**, 115–119, 2000; T. M. Davis and C. H. Lineweaver, *Pub. Astron. Soc. Austral.* **21**, 97–109, 2004; L. A. Barnes, M. J. Francis, J. Berian James, and G. F. Lewis, *Mon. Not. R. Astron. Soc.* **373**, 382–390, 2006; M. J. Chodorowski, *Mon. Not. R. Astron. Soc.* **378**, 239–224, 2007; G. F. Lewis, M. J. Francis, L. A. Barnes, and J. Berian James, *Mon. Not. R. Astron. Soc.* **381**, L50–L54, 2007; R. J. Cook and M. S. Burns, *Am. J. Phys.* **77** (1), 59–66, 2009.
- [8] N. V. Budko, *submitted to Phys. Rev. A*, see also: arXiv:0906.0495, 2009.
- [9] E. Schrödinger, *Physica* **VI**, 9, 899–912, 1939.
- [10] J. Plebanski, *Phys. Rev.* **118**, 5, 1396–1408, 1960.
- [11] F. de Felice, *Gen. Rel. Grav.* **2**, 4, 347–357, 1971.
- [12] M. Bahram, *Phys. Rev. D* **8**, 12, 4297–4302, 1973.
- [13] J. E. Tamm, *J. Russ. Phys.-Chem. Soc.* **56**, 2-3, 284, 1924.
- [14] U. Leonhardt and T. G. Philbin, *New J. Phys.* **8**, 247, 2006.